

2014-2015 School Nurse Guide
Wyoming School Immunization Requirements (Kindergarten – Grade 12)
 (Updated 06/2014)

Minimum Immunization Requirements: 1. Parents/guardians must provide proof of immunization or provide documentation of medical or religious exemption from mandatory immunizations; 2. A vaccine administered up to four (4) days prior to the minimum age/interval for that vaccine is considered compliant with that vaccine administration schedule.

Vaccine	Kindergarten - Grade 6	Grade 7 - 12	Footnotes
Diphtheria/Tetanus/acellular Pertussis (DTaP) or Diphtheria/Tetanus/Pertussis (DTP) or Diphtheria/Tetanus (DT)	5 doses ¹	Complete Series (i.e. Td if >age 7)	1. If a fourth (4th) dose of DTaP vaccine was administered on or after a child's fourth (4th) birthday, and at least six (6) months has passed since the third (3rd) dose, a fifth (5th) dose is not required.
Haemophilus influenzae type b (Hib)	"Complete Valid Series" (i.e. 1-4 doses*)	n/a	*The number of primary doses of Hib vaccine is determined by vaccine product and age the series begins. If a child has not received the completed series of Hib vaccine and is younger than 60 months (5 years) of age, the child should receive one (1) dose of the Hib vaccine. Hib vaccine is not required for children age 5 years and older.
Hepatitis B (Hep B)	3 doses	3 doses	
Measles/Mumps/Rubella (MMR)	2 doses	2 doses	
Polio (IPV)	3-4 doses ²	3-4 doses	2. Children who receive three (3) doses of IPV before the fourth (4th) birthday should receive a fourth (4th) dose before or at school entry. The fourth (4th) dose is not needed if the third (3rd) dose is given on or after the fourth (4th) birthday. If all four (4) doses are given after six (6) weeks of age and are all separated by at least four (4) weeks, a fifth (5th) dose is not needed, even if the fourth (4th) dose was administered before four (4) years of age.
Tetanus/diphtheria/acellular pertussis (Tdap) ⁵	n/a	1 dose**	**Children eleven (11) years of age and older should receive the Tdap vaccine. Children with a medical contraindication to the pertussis (whooping cough) antigen should receive Td instead of the Tdap vaccine. Tdap vaccine may be administered to a child regardless of the interval since the last tetanus- or diphtheria-toxoid containing vaccine. During the 2014-2015 school year, any student in grades eleven (11) through 12 is considered exempt with the immunization requirements for Tdap, according to when the new school rules became effective on 08/17/10.
Varicella (chickenpox)	2 doses (or documented history of disease)***	2 doses (or documented history of disease)***	***If a child has had chickenpox, the parent, guardian, school nurse, or physician must provide a written statement for the student's permanent school record. A student who has been diagnosed with a history of herpes zoster (i.e. Shingles) is considered as having evidence of immunity to Varicella.

Resources:

- WY Immunization School Rules & this Guide:: <http://www.health.wyo.gov/familyhealth/immunization/SchoolResourcesSchoolImmRecs.html>
- School Nurse Immunization Liaison: Diana L. Martin, RN, MS, BSW, CPHQ, Clinical Quality & Compliance Specialist, WY Dept. of Health - Immunization Program , Telephone: (307) 787-6052, E-mail: diana.Martin@wyo.gov